

## Using Fentanyl Test Strips (FTS)



If you use the test strips provided, you acknowledge and accept all responsibility and liability for possessing the test strips.

A wide variety of substances can be laced with fentanyl. While it is advisable to test substances for fentanyl, there are no well-researched, established recommendations on how to test drugs with test strips.

There is a possibility that the test strips may not indicate the presence of fentanyl and they are not guaranteed to detect all forms of fentanyl. You could overdose even if the strip says there is no fentanyl in your heroin, cocaine, or other substances. If you use the FTS provided, you accept all responsibility and liability for any injury or death that could occur after ingesting any substance, whether they have been tested, or not tested for fentanyl.

### Testing for Fentanyl


- Prepare shot and draw it into a syringe leaving several drops in the cooker. Do not shoot up yet!
- Set the loaded syringe aside and put clean water into the cooker.
- Stir the mixture.
- Holding a test strip by the blue end, insert it into the solution (don't go past the blue line) and leave it there for about 15 seconds.
- Wait a full minute for the test strip to process.
- If using pills, crush them first, and then the powdery residue can be tested by mixing with a bit of water and following the directions above.

One Red Line = Positive for Fentanyl  
Two Red Lines = Negative for Fentanyl

Knowing fentanyl is present in your substance gives you some options, you can:

- choose not to use it;
- start with a smaller amount;
- go slow while using.

Always have Narcan available and call 911 in an overdose.  
An overdose required medical attention!



Scan for  
Free  
Narcan



**MAHONING COUNTY  
PUBLIC HEALTH**  
PREVENT · PROMOTE · PROTECT